



FOR ANY AND EVERY LOOK



CLOSEST TO NATURAL HA

The original HA filler with minimal modification^{1,2}



HIGH PERFORMANCE – LESS PRODUCT FOR GREAT EFFECT THAT LASTS³⁻⁵

- Less product needed to achieve the desired result^{3,4}
- Results that last up to 24 months, with one retreatment depending on the product used and area treated $^{\rm 5}$



EXPERTS IN FACIAL DYNAMICS

Clinically proven to have positive effects on the naturalness of dynamic facial expressions^{6,7}

PROVEN SAFETY



Unmatched 26 years of proven safety in nearly 407,000 patients with diverse skin colors and types^{**+5,7-10}

 Restylane® products have demonstrated a potentially lower risk of late-onset complications compared with Juvéderm® products¹⁰



HIGH PATIENT SATISFACTION ACROSS 26 YEARS**11-13

26 years of experience and trusted with more than 55 million treatments worldwide¹¹

Your expertise, meets ours. With the world's most diverse range of fillers and skinboosters.^{§14-24} Each and every one individually engineered so you can bring bespoke looks to life with precision.^{25,26} Matching your mastery, with their vision.

FOR ANY AND EVERY LOOK

WITH DIFFERENT TECHNOLOGIES SUITED FOR EACH TREATMENT AREA. YOU CAN CREATE A LOOK TAILORED TO EACH PATIENT.



Restylane



- A second treatment (retreatment) was performed at 12 months in the same areas with the same products.⁵ Touch-up was allowed 1 month after both treatments.⁵ Restylane^{*} injectable gel (also known as Restylane^{*}) was first approved for marketing and sale in September 1996 in the EU.¹¹ Based on a Restylane^{*} publications analysis conducted in June 2021.⁹ Sources include a published post-marketing safety study of 638 patients and a retrospective analysis of adverse event reports encompassing 406,000 patients.⁹ In a recent post-marketing safety surveillance study using the MAUDE database to assess delayed (zl4 days post-treatment) AEs of interest (inflammatory and noninflammatory nodules, hypersensitivity, granulomas), Restylane^{*} Refyne^{**} and Restylane^{**} Defyne^{**} both had fewer combined total events of interest vs. Juvéderm^{**} Volbella^{**} and Juvéderm^{**} Voluma^{***}.¹⁰ The Restylane^{**} product family has a broader range in G^{*} (elastic modulus) than Juvéderm^{**}, Belotero^{**}, and Teosyal^{****}.^{14,15} G^{*} is a measure of gel firmness (resistance to deformation), which can be used to understand how a gel¹⁵ physical properties relate to its in vivo nerformance¹.
- ş (resistance to deformation), which can be used to understand how a gel's physical properties relate to its in vivo performance.

AE, adverse event; HA, hyaluronic acid; MAUDE, Manufacturer and User Facility Device Experience.

- References:
 Öhrlund Å. J Cosmet Dermatol Sci Applic 2018; 8(2): 47-54.
 Kablik J et al. J Dermatol Surg 2009; 35: 302-312.
 Hilton S et al. Dermatol Surg 2018; 44: 261-269.
 Jones D et al. J Drugs Dermatol 2021; 20: 949-956.
 Huang S and Tsai T. J Drugs Dermatol 2020; 19(9): 836-842.
 Philipp-Dormston W et al. Dermatol Surg 2018; 44: 826-832.
 Percec I et al. Plast Reconstr Surg 2002; 145(2): 295e-305e.
 Kerscher M et al. Dermatol Surg 2008; 34: 720-726.
 Galderma. Data on File. MA-48087.
 Galderma. Data on File. MA-48080.
 Galderma. Data on File. MA-48544.
 Andriopoulos B et al. Poster presented at Aesthetic & Anti-aging Medicine World Congress. 4-6 April 2019. Monte Carlo, Monaco.
- **14.** Galderma. Data on File. MA-43049 **15.** Galderma. Data on file. MA-34483.

- Galderma. Data on file. MA-34483.
 Restylane* EU IFU.
 Restylane* Volyme™ EU IFU.
 Restylane* Volyme™ EU IFU.
 Restylane* Kysse™ EU IFU.
 Restylane* Skinboosters™ Vital EU IFU.
 Restylane* Skinboosters™ Vital Light EU IFU.
 Restylane* Skinboosters™ Vital Light EU IFU.
 Restylane* Lyft™ EU IFU.
 Restylane* Skinboosters™ Vital Light EU IFU.
 Restylane* Lyft™ EU IFU.
 Restylane* Skinboosters™ Vital Light EU IFU.
 Restylane* Lyft™ EU IFU.
 Restylane* Lyft™ EU IFU.
 Restylane* Lyft™ EU IFU.
 Restylane* Lyft™ EU IFU.